

King's Research Portal

DOI:

[10.1016/j.ecoser.2018.03.018](https://doi.org/10.1016/j.ecoser.2018.03.018)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Kim, Y-S., Latifah, S., Afifi, M., Mulligan, M., Burke, S., Fisher, L., Siwicka, E., Remoundou, K., Christie, M., Masek Lopez, S., & Jenness, J. (2018). Managing forests for global and local ecosystem services: A case study of carbon, water and livelihoods from eastern Indonesia. *Ecosystem Services*, 31(Part A), 153-168.
<https://doi.org/10.1016/j.ecoser.2018.03.018>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Full title: Managing forests for global and local ecosystem services: A case study of carbon, water and livelihoods from eastern Indonesia

Highlights

- Trade-offs/synergies exist among ecosystem services in different scales.
- Forest carbon projects can have negative effects on watershed services.
- Watershed services from forests are very important to local communities.
- Forest carbon projects must consider further details within a land use class.

Full title: Managing forests for global and local ecosystem services: A case study of carbon, water and livelihoods from eastern Indonesia

Abstract

Despite a recent increase of interest in global payment for ecosystem services (PES) mechanisms, there has been little comprehensive assessment of PES impacts on ecosystem services (ESs) at smaller scales. Better understanding of localized impacts of global PES can help balance ES deliveries for global benefits with those for meeting landscape and local level needs. Using a case study from eastern Indonesia, we assessed trade-offs and potential synergies between global PES (e.g. REDD+ for forest carbon) and landscape level ESs (e.g., water quantity, quality, regulation) and local ESs (e.g. forest products for food, energy, livelihoods). Realistic land use change scenarios and potential carbon credits were estimated based on historical land use changes and in-depth interviews with stakeholders. We applied a process-based hydrologic model to estimate changes in watershed services due to land use changes. Finally, local community's forest uses were surveyed to understand locally realized ESs. The results show empirical evidence that, without careful consideration of local impacts, a PES mechanism to protect global ESs can have negative consequences for local ecosystem services. We present management alternatives designed to maximize positive synergies between different ESs at varying scales.

Keywords: ecosystem services; carbon; REDD+; watershed services; livelihoods; Indonesia,

1 Introduction

Globally, tropical forests account for approximately 25% of all terrestrial carbon (Bonan, 2008). Deforestation is the largest source of carbon emissions from tropical developing countries (Pan et al. 2011). The 2015 UN climate change conference in Paris reconfirmed the importance of forests in global climate regulation. The agreement explicitly included the REDD+ mechanism¹ as part of the global climate regime, where tropical and sub-tropical countries could receive both public and private funding for reducing carbon emissions and conserving standing forests. Indonesia has the third largest tropical forest in the world, with one of the world's fastest rates of deforestation at more than 1,000 km² of forests (476 km² of primary forest) lost per year between 2000-2012 (Hansen et al., 2013; Margono et al., 2014). Indonesia has emerged as the major beneficiary of global negotiations to mitigate climate change through improved forest management (Simula, 2010). It has received the largest portion of REDD+ readiness commitments from the public sector (\$757 million out of \$2.8 billion total committed and dispersed from 2009 to 2014; Goldstein et al., 2015). In the private sector, carbon credits from protecting Indonesia's forests was 5.5% of all voluntary carbon transactions in 2015 (Hamrick and Goldstein, 2016).

Offering financial incentives for tropical developing countries to reduce deforestation and forest degradation can be a win-win-win solution for climate mitigation, ecosystem conservation and poverty alleviation (Pistorius, 2012). However, many previous studies have warned that international intervention in the form of Payments for Ecosystem Services (PES) can exacerbate internal social problems (Blom et al., 2010; Wunder, 2008). Failure to include consideration for local uses of resources in global PES design can

¹ Reducing Emissions from Deforestation and Forest Degradation (REDD+) is an effort to offer financial incentives for developing countries to reduce emissions from forested lands. REDD+ projects include activities for (a) reducing emissions from deforestation, (b) reducing emissions from forest degradation, (c) while recognizing the role of conservation of forest carbon stocks, (d), sustainable management of forests, and (e) enhancement of forest carbon stocks (UN-REDD programme, 2017).

undermine rights of indigenous and local communities, exacerbate food and water insecurity (UN-REDD programme, 2017; Fazey et al, 2010), diminish ecological integrity and equity (Motel et al., 2009), and result in less than optimal outcomes for the ecosystem service targeted (Enrici and Hubacek, 2016; Skutsch et al. 2011). Despite a recent increase of interest in global PES mechanisms, there has been little comprehensive assessment of their impacts on localized ecosystem services (ESs) and livelihoods. Better understanding of the localized impacts is needed to find ways of balancing provision of an ES that provides benefits at the global scale, while meeting local needs for water, food, energy and livelihoods. Using a case study from eastern Indonesia, we present a detailed assessment of trade-offs and potential synergies among global ES (forest carbon), landscape-level regulating services (e.g. water) and localized provisioning services (e.g., forest products for food and energy). Specific research questions are: 1) what are realistic land management scenarios to recover forest area lost and improve forest conditions?; 2) how do these scenarios affect global, landscape and local ES provisions?; 3) how do global modelling results compare with local perception in assessments of ecosystem service change; 4) what are the management alternatives to maximize positive synergies among provisions of different ESs at varying scales?

2 Literature review: Ecosystem Services trade-offs and synergies

The Millennium Ecosystem Assessment (MA 2005) placed the term “ecosystem services” firmly in the policy agenda (MA 2005; Gómez-Baggethun et al., 2010). Since then, many have advocated the urgent need to incorporate sustainable provisioning of ESs into policies and planning for managing landscapes (e.g., Daily et al. 2009; de Groot et al, 2010). However, the flows of ESs are determined not only by ecosystem functions and processes (ES supply), but also by demands from various human actors (ES demand) in multiple-scales (Figure 1). Mouchet et al. (2014) advanced a typology to understand ES trade-offs by merging ecological and socio-economic considerations found in previous studies. Spatial and time lags of ESs (spatial and temporal trade-offs) can occur in both supply and demand sides, in terms of production and delivery (Rodriguez et al. 2006) and benefits and costs (TEEB, 2010). Also targeting one ES can affect other ESs positively or negatively

(among ESs synergies or trade-offs), and resilience of the ecosystem as a whole (reversible trade-off), as well as who “losers” and “winners” are among ES beneficiaries (beneficiaries trade-off) (Mouchet et al. 2014).

<Figure 1>

The forces of globalization are intensifying interactions among ES demand and supply over distances and cross-scales (Cash et al. 2006; Liu et al. 2015). Managing ESs and anticipating changes in their spatial, temporal and societal distributions are increasingly difficult as local events (e.g. land use change in tropics) can have global consequences (e.g. climate change) (e.g. Bruckner et al. 2015; Meyfroidt et al. 2013; Lambin et al. 2011; Seto et al. 2012). Spatially distributed beneficiaries of different ESs vary also in their social and economic status, which affect their ability to influence decision-making process (TEEB 2010). There have been several studies that addressed the spatial scale of managing ES (e.g., Hein et al., 2006; Willemen et al., 2012 – both in the Netherlands) and presented empirical evidence of trade-offs and synergies of different ES deliveries (e.g. González-Esquivel et al. 2015; Grossman 2015; Haines-Young et al., 2012; Maes et al. 2012; Mastrangelo and Laterra 2015; Mora et al. 2015; Turner et al. 2014 – in Europe and Latin America), However, those most affected by global PES, such as REDD+, are in tropical developing countries often lacking technical capacity for data collection, analysis and management (Goetz et al. 2014). With the growing significance of global carbon governance (Bierman, 2010), there is a critical need to understand how the economic and political scale of decision-making affects ESs at different scales. We chose three groups of ESs at global, landscape (watershed level) and local community scales to contribute to our current understanding about ES associations and potential effects of global PES schemes.

3 Methods

3.1 Study area

The case study area is Lombok island in Nusa Tenggara Barat (NTB) province, located in eastern Indonesia (Figure 2). According to a recent analysis of Landsat images, the forested area of Lombok decreased 28.6% from 1990 to 2010 (Bae et al., 2014). By comparison,

100 Indonesia's national average forest loss is 20.3% during the same period (FAO, 2010).
101 Lombok is also one of the most densely populated and impoverished areas in Indonesia.
102 Seventy percent of the population of NTB province lives in Lombok, although the island
103 only constitutes a quarter of the total land area of the province (708 persons/km²,
104 compared to 237 persons/km² for NTB and 132 persons/km² nationally, as of 2014, BPS-
105 NTB, 2015). Economic opportunities are limited to agriculture (24% of Gross Domestic
106 Product (GDP) and 43% of employment of the province) and the mining and quarrying
107 sector (15% of GDP and 1.8% employment) (as of 2014, BPS-NTB, 2015). NTB is among
108 the poorest provinces of Indonesia, based on the Human Development Index (HDI), a
109 metric that combines average life expectancy, education level, and *per capita* income (65.19
110 compared to the national average 69.55 as of 2015, BPS, 2016).

111 <Figure 2>

112 Although forestry is a relatively small contributor to the wider economy of NTB (0.1% of
113 GDP as of 2014, BPS-NTB, 2015), the forests in the northern part of the island, surrounding
114 the Rinjani volcano complex, are an important source of subsistence and income to local
115 communities. The forest also represents an important watershed, providing municipal
116 water for the city of Mataram and irrigation for the major rice production regions
117 throughout Lombok Island. The development of a program of payment for watershed
118 services between municipal rate-payers and forest margin communities is one of the very
119 first examples of PES systems in Indonesia (Diswandi, 2017; Pirard 2012; Prasetyo *et al.*,
120 2009). The program supports forestry or agroforestry projects proposed by community
121 groups with funds collected from the downstream city's water use fees. A multi-
122 stakeholder group (IMP, *Institusi Multi-Pihak*) consisting of representatives from the
123 World Wildlife Fund, the district forest service, a local university, a mineral water
124 company, the district government and Mount Rinjani National Park, selects and distributes
125 funds for selected projects (Diswandi, 2017; Schweizer et al. 2016; Pirard, 2012).

3.2 Research approach

To assess the potential impacts of different land use change scenarios on ESs at different scales, we first identified alternative forest management scenarios that can be adopted by a future carbon PES scheme in Lombok. We then assessed the carbon, water and locally important services for food, energy and livelihoods impacts of these PES scenarios.

3.2.1 Forest management scenarios

Forest carbon projects are designed to provide incentives to protect forests for the value of their standing carbon. Estimating carbon credits is essential for establishing the economic value of forest carbon projects. It includes two components: land-use and land-cover changes and the associated changes in carbon stock (VCS, 2012).

Future forest management scenarios were developed based on analysis of historical changes in land-use and land-cover, along with analysis of drivers of deforestation and forest degradation in the area. Detail of these changes have been reported in Bae et al., 2014 and Kim et al., 2016. Table 1 shows the changes in deforestation patterns in three 5-year periods (1995-2000; 2000-2005; 2005-2010). Land use classes² following deforestation were projected based on the satellite imagery footprint of the most recent historical land cover pattern (2005-2010). We focus on the area around the Rinjani volcano complex, where the majority of Lombok's remaining forests are located.

<Table 1>

² Primary forest in this study is defined as mature or intact forest, where standing stock has reached stability. The forest is generally of native tree species, there are no clear indications of human activities, and the ecological processes are not significantly disturbed. Secondary forest is regenerated forest that has been disturbed by human activities or natural disasters. Secondary forest may include a natural forest with timber extraction, retaining artificial gaps in the canopy to 50-60%. This kind of forest includes agroforestry and community forests. Shrubland refers to land with woody vegetation where the dominant woody elements are shrubs, bushes and young generation trees, generally less than 5m in height. The latter appears usually after forest clear-cutting activities without crop cultivation. (Source: Bae et al. 2014).

When the Suharto regime fell in 1998, this socio-political shift caused an abrupt interruption of central government control of forest lands that encouraged massive forest encroachment that was common throughout Indonesia at the time (e.g., Resosudarmo, 2004). Figure 3 graphically illustrates the deforestation patterns during the three 5-year periods studied. Between 1995 and 2000, land use changes were driven by conversion of primary and secondary forests to shrubland, indicating no immediate cultivation after clearing of forest lands. After 2000, deforestation of primary forests decreased and some shrubland transitioned back to secondary forest. However, deforestation of secondary forest continued and secondary forest and shrubland are now being cultivated for dryland agriculture.

<Figure 3>

In addition to examining the historical patterns of land use changes, we conducted a series of in-depth interviews (January 2015) with key informants from provincial and local government forest agencies, as well as international and local NGOs, to better understand the varied contexts of forest management. Based on this information, we develop three land-use change scenarios that represent a range of possible reforestation and restoration outcomes. These scenarios are reported in Section 4.1.

3.2.2 Carbon assessment

To estimate the impacts of the projected future land use changes on carbon stocks, we used the area-weighted average of carbon stock for each carbon pool for forest and shrubland, based on field inventory (Table 2). The estimated changes of carbon stock are based only on land use class change in each scenario and do not incorporate other variations within land use classes. For all other land uses, the carbon stocks were assumed to retain the level of soil carbon in shrubland³.

³ For carbon stock change, Verified Carbon Standard (VCS) guidelines state that the REDD+-related projects should account for the following carbon pools: above-ground living biomass of trees and non-trees, and wood products if harvested timbers are utilized to make long-lived wood products. Measuring and monitoring

<Table 2>

3.2.3 Hydrological modelling

We utilized a process-based hydrologic model, WaterWorld V 3.31, to project the hydrological impacts of the land-use change scenarios. WaterWorld is a spatially explicit, globally applicable model for calculating monthly water balance, runoff, water quality (including agricultural pollutants and soil erosion) and their spatial distributions under baseline and alternative land use change scenarios (Mulligan, 2013). WaterWorld V 3.31 uses globally available data sets from remote sensing, along with limited *in situ* precipitation data to reveal how forest restoration can affect water provisioning and regulating services (Mulligan 2013). WaterWorld V 3.31 calculates water balance as a sum of wind driven rainfall, fog and snowmelt (not applicable in this case) minus actual evapotranspiration. Water infiltrates according to regional infiltration capacities (Gleeson et al., 2011), mediated by slope gradient and tree cover (lower gradient and greater tree cover lead to higher infiltration rates within the geology-controlled regional limits). Infiltration is calculated based on global permeability data using the lithology developed by Gleeson et al. (2011). The infiltration model takes the mean soil-conditioned hydraulic conductivity as the infiltration rate and increases it towards one standard deviation higher than the mean in each pixel as tree cover increases and slope decreases. Higher tree cover encourages infiltration, shallower slopes provide greater opportunity for it to occur. Infiltration is also limited by available porosity and declines in a linear fashion as the soil store fills. Infiltrated water joins subsurface base flow and travels much more slowly to streams than water running over the land surface. Infiltrated water flows downslope along subsurface flow lines dictated by surface topography and at rates dictated by the local infiltration rates of the soil that water is passing through. Infiltrated water may re-emerge as surface runoff anywhere downslope where soil conditions (subsurface flow rates) or water conditions (volume of water in relation to soil thickness mediated storage capacity)

other carbon pools, such as living below-ground biomass and dead organic matter, are optional or not required.

dictate. This tends to occur most at the base of hillslopes and in channels where regolith thickness is less and thus water emerges at the surface, as baseflow. There is no separate deep groundwater model, WaterWorld models all subsurface moisture as a single per pixel unit. Tree cover also increases the rate of evapotranspiration and the rate of interception of fog, where it occurs. The model was applied to the current conditions in Lombok to produce information on the current hydrological ESs and also model their changes under different land use change scenarios. We also assessed local perception of watershed services linked with forest conditions through focus group discussions (FGD) and survey.

3.2.4 Locally important ecosystem services for food, energy and livelihoods

To understand how local community members utilize and benefit from forest ecosystem services, in-person surveys were conducted at four locations (Figure 4). Survey locations were selected based on their proximity to forests with different designated functions, forest governance status, and permitted activities.

State forests in Indonesia are classified into three designated functional categories (ROI, 1999)⁴: 'Production Forest' for providing forest products; 'Protection Forest' for ecosystem protection, such as watershed and soil conservation; and 'Conservation Forest' for protecting biodiversity and ecosystem conservation. Production and Protection Forests in NTB province are managed by Forest Management Units (*Kesatuan Pengelolaan Hutan*, or KPH) that were created by the central government but are more or less decentralized (See Kim et al., 2016 and Sahide et al., 2016 for more complete information on the Forest Management Units). Conservation Forest is directly managed by the National Park (i.e. Conservation Forest Management Unit) under the central government authority. We selected one community adjacent to Production Forest (A), one near Protection Forest (B), and one near Conservation Forest (C), i.e., near the Rinjani National Park (Figure 4).

⁴ Indonesian Law Number 41/1999 distinguishes "forest" as an ecosystem dominated by trees and "forest area" defined as a particular area designated by the government. Thus, these administrative designations may not necessarily represent actual forest cover and particular forest conditions (Bae et al. 2014)

219 <Figure 4>

220 We also included an additional community near a Protection Forest that recently gained
221 official recognition as “Community Forest” (*Hutan Kemasyarakatan*, or HKm) (D).
222 Community Forest is one of the legal mechanisms that communities can use to gain
223 recognition for their usufruct rights (ROI, 2007). However, the legal process of establishing
224 HKm is complicated, involving both local and central government agencies, and it can take
225 years to gain formal approval (Intarini et al., 2015), which explains why less than 1% of
226 Indonesia’s forests are managed by communities with HKm status (Stevens et al. 2014)⁵.
227 This particular community gained HKm status through intense facilitation supported by an
228 international NGO (Flora and Fauna International) that initiated a REDD+ demonstration
229 project in the area.

230 The various forest designations offer alternative levels of forest protection. As such, they
231 differ in terms of the activities that local people are permitted to undertake in the forest.
232 Table 3 provides a summary of permitted activities by forest designation.

233 <Table 3>

234 We conducted surveys across locations A, B, C, and D (January 2015) to assess the
235 importance that community members attach to local forest ESs across the four locations. A
236 list of locally important forest ESs was drawn up, following scoping focus group discussions
237 with community members and local stakeholders. These services were then grouped into
238 three groups of provisioning services and one regulating service:

- 239 • Naturally occurring non-timber forest products (NTFP), such as bamboo, honey and
240 cattle feed;
- 241 • Agroforest products, such as various fruits and cash crops (e.g., coffee and cacao);
- 242 • Timber forest products, including fuelwood; and

⁵ The government of Indonesia declared a plan to dramatically increase community control of forests from 1.4 million hectares in 2014 up to 12.7 million hectares by 2019 and is currently identifying the areas suitable for community forests (Indonesia National Planning & Development Agency, 2015)

- Water regulation services.

Although cultural services of forests were also identified to be significant to these forest margin communities, it is difficult to measure those services and link them to forest conditions. Thus they were not explicitly investigated in our study. The survey questionnaire comprised five sections. First, we collected background information on the respondents, including their proximity to the forest. Next, we asked a general question on the extent to which the services they obtain from the forest sustains their needs and how this has changed over the past 5 years. The third and fourth sections respectively collected detailed information on the levels of consumption of provisioning and regulating services. Finally, we collected information on respondent's preferences for alternative future forest management options. The surveys were administered in-person by (trained) local enumerators, who conducted the surveys in the respondent's home in the local language. A sampling frame was developed for identifying respondents following consultation with community leaders and aimed to obtain a representative sample of community members. Survey data was analyzed separately for the four locations. After analyzing the data, we held a workshop with community members in each location to share our findings, elicit feedback on our preliminary results, and explore possible future options to more effectively manage the forests (March 2016).

4 Results

4.1 Land use change scenarios

Three future (30-year projection) land use change scenarios were developed based on spatial data on recent land use changes (2005-2010), combined with current forest management plans obtained from key informant interviews (January 2015). The scenarios included a Business-As-Usual scenario and two management scenarios aimed at improving forest condition.

268 4.1.1 Business-As-Usual (BAU) scenario

269 There has been little decrease of primary forests in the study area since 2000, although
270 secondary forest and shrubland have changed to other land uses, primarily dryland
271 agriculture. Under this scenario, these current trends in land use change would continue
272 unabated, resulting in ~10% of currently forested land being converted to dryland
273 agriculture. We used the latest available land-use data (2010) as the starting point for our
274 simulations. The projected land use changes for the next 10 and 30 years are shown in
275 Table 4

276 <Table 4>

277 4.1.2 Community Partnership (CP) scenario

278 Forest Management Units (KPHs) in Lombok currently use a spatial planning approach, in
279 which the remaining primary forests are defined as core protected zones, and surrounding
280 secondary forests are designated for community use. The agencies are developing
281 programs to assure *de facto* usufruct rights for communities and allow agroforestry
282 development through partnership agreements (*kemitraan*) in the secondary forest (Jang
283 and Bae, 2014). The optimistic, yet realistic, scenario would be that this program will
284 succeed at buffering encroachment into the primary forest, and the partnership
285 agreements will expand to all forests around Mount Rinjani managed by KPHs. The
286 resulting land use changes would increase the area of secondary forests to the 1995 level
287 (i.e. before the period of rapid deforestation) with 50% of forest restoration occurring in
288 the first 10 years. In this scenario, secondary forests would include well-managed
289 agroforestry areas with forest cover converted from shrubland (32% increase of total
290 forests in 30 years), while the area of primary forests would remain unchanged (Table 5).

291 <Table 5>

4.1.3. Forest Restoration (FR) scenario

This scenario presents the realistic upper limit of a reforestation scenario. It would require an intervention, for example a REDD+-type carbon project, that would lead to restoring all Lombok's forests to the 1995 levels with 50% of forest restoration occurring in the first 10 years. The resulting land use changes would include 7% increase of primary forest and 56% increase of total forest in 30 years (Table 6).

<Table 6>

4.2 Changes in carbon stock and potential carbon market values

Table 7 shows land use changes under two scenarios compared to the BAU scenario, as well as resulting total carbon stock changes. For example, secondary forests in Lombok, which contain an average of 181.1 metric tons of carbon per ha, are projected to increase by 24,060 ha in 10 years under CP scenario (from 65,462 ha under BAU to 89,522 ha under CP scenario). After combining changes in carbon stock with all land uses, total carbon stock under CP scenario would be a 4.0 million metric tCO₂e increase for the first 10-year period, and a 6.9 million metric tCO₂e over the thirty year project period. FR scenario will result in increase of 4.3 million metric tCO₂e from BAU scenario REL for first 10 years and 7.6 million metric tCO₂e over the 30 year project period.

<Table 7>

Carbon price (USD/ metric tCO₂e) in voluntary carbon market varies by sources, although it is commonly higher for forest carbon. REDD+ projects for avoided planned deforestation (\$1.9) and avoided unplanned deforestation⁶ (\$5.5) generally resulted in forest carbon offsets whose values were lower than those from sustainable agriculture/agroforestry

⁶ Carbon credits from REDD+ projects are based on different forms of avoided emission from planned (i.e. legally authorized and documented for conversion) and unplanned deforestation, as well as forest degradation (i.e. canopy cover remaining above the threshold for definition of forest and no change in land use).

(\$7.4), tree planting (\$8.9) and improved forest management (\$9.8) projects (average prices per metric tCO₂e in 2014 from Goldstein et al., 2015). Even at the lower end of carbon price (\$5) and emission reduction, we can expect at least \$35 million of expected value generated for a 30-year forest carbon project in Lombok (Table 8). However, this amount indicates the carbon credit potential, not necessarily the actual payments required to start a project.

<Table 8>

4.3 Hydrological modelling results

WaterWorld V3.31 results predicted that CP and FR scenarios would result in decreased local annual water balance and runoff in most locations in Lombok due to increased evapotranspiration from tree cover. Figure 5 shows the changes in average surface water runoff and water balance under CP and FR scenarios. The differences between catchments reflect differences in the amount of tree cover change as well as the effects of varying fog frequency, rainfall totals and slope.

<Figure 5>

The WaterWorld metric for water quality is termed the human footprint on water quality (Mulligan, 2010; Mulligan, 2013) and indicates the impact of upstream land use on downstream water quality as a percent of water that fell as rain on human impacted land uses. Water quality was predicted to increase in the afforested areas because of reduced agricultural inputs, but reduced runoff through greater evapotranspiration can also translate to concentrated pollutants downstream from the remaining agricultural lands. Since most populations are at lower elevations (e.g. residents in the city of Mataram. For the location, see Figure 1) and most forest are at higher elevations, this can mean a minimal or negative effects from increasing forest cover on water quality to downstream beneficiaries. Moreover, although increased infiltration does lead to a greater fraction of water as subsurface flow, WaterWorld V3.31 shows the impact of reduced water balance is greater so dry season flows decrease as tree cover increases in this region. Overall, the

water modeling showed no net benefits from recovering tree cover in terms of water supply and water quality downstream, except locally at a few remote very cloudy sites.

4.4 Local perceptions of forest ESs

To assess potential impacts of future land use change scenarios on provisioning services that sustain food, energy and livelihoods of local communities, we surveyed 408 individuals across the four forest locations. During the surveys, respondents were asked to report on their household's level of consumption of forest ESs obtained from the forest (NTFPs, agroforest products, and timber products), and their perceived market values of these ecosystem services (Section 4.4.1). We also asked respondents to indicate what services they would like to see being enhanced through future forest management actions (Section 4.4.2).

4.4.1 Locally important provisioning services from forests

The majority (80%) of respondents reported that their household utilizes some forest ESs (Table 9). The community near the Protection forest (B) reported highest level of use (98% of respondents), followed by A near Production forest (86%), C near Conservation forest (81%), D Community forest (53%). Agroforest products were utilized most widely (69%), while smaller portions of respondents reported utilization of Natural NTFP (49%) and Timber (47%). The specific forest products utilized vary by locations: coffee (67%), banana (56%) and fern (49%) were most popular in A community; jackfruit (86%) and banana (82%) in the B community; fern (69%) and forage (58%) in C community; and coffee (35%) and Jackfruit (34%) in D Community forest. Fuelwood collection was higher in A near the Protection forest (79%), compared to other areas around where one-third of respondents reported collection. These variations are due to differences in permitted activities across different forest designations (See Table 3), as well as ease of access to markets and other socio-economic variables. For example, a previous study showed that domestic energy needs can be often met by deadwoods and branches collected in household gardens and fuelwood extraction from forests is highly correlated with opportunity to sell fuelwoods (Lee et al. 2015).

We also explored the economic value of the products collected from different locations. To calculate these values, reported volumes collected were multiplied by reported prices. When the price was missing but the respondent reported some level of extraction, the mean price was used. To get a conservative estimate of the values and avoid outliers skewing the data, we removed the top and bottom 10% of the value distribution. Average overall values of forest ESs utilized per household per year were highest in the Production forest (\$141), followed by Community forest (\$116), Protection Forest (\$85) and Conservation forest (\$46).

Table 9 provides further detail of the distribution of values by ESs by location. Highest values were found for Palm (\$918 for 6% of Community forest users), Coffee (\$262 for 67% of Production forest users and \$64 for 35% of Community forest) and Durian (\$81 for 13% of Community forest users and \$75 for 33% of Production forest users). Timber products were largely restricted to fuelwood with relatively low value (\$4/household/yr). Forest products most likely to be consumed by the household are: melinjo (94%), forage (91%), jackfruit (88%), taro (83%) and fern (83%), while cacao (92%) and palm (83%) were the products most likely to be sold. Our findings demonstrate that there was a significant variability in terms of forest uses by communities.

<Table 9>

4.4.2 Perceived importance of forest ESs

We asked respondents to indicate which services they would like to see improved by future forest management plans. Both water regulation (91% of respondents) and agroforest products (81%) were considered to be important by most respondents; a finding that is consistent across all four forest locations (Table 10). The over-riding importance placed on water regulation can be illustrated by a comment made by one respondent “[Other ecosystem services] are what we need to live, but water is life”. The higher importance ranking of agroforest products may be explained by the fact that more people used and obtained higher values of services from agroforest products than the other forest ESs categories (Table 9). Natural NTFP (40%) and timber (27%) were considered to be less

important. However, there were significant differences between locations in terms of the importance of these services. Natural NTFPs were considered important (67%) in the Conservation forests, while timber resources were considered important (76%) in the production forest. These differences in preferences reflect the activities that are permitted in the different types of forest. Analysis of the socioeconomic characteristics of respondents indicated that, generally, there was little difference between the socio-economics of the people living in the different forests.

<Table 10>

5 Discussion

5.1 Forest management, PES and the delivery of global and local services.

In this research, we explored the potential impacts of alternative land use change scenarios on ecosystem services across different scales from global to landscape and local levels. Our analysis identified two scenarios: a community partnership (CP) scenario which largely focused on increasing the area of secondary forest; and a forest restoration (FR) scenario which increased the area of both secondary and primary forest. In terms of global ES, it is clear that both of these scenarios can generate significant global carbon benefits: over a 30-year period the CP scenario was estimated to generate between \$35 million to \$69 million in carbon values, while the FR scenario would generate between \$38 million and \$76 million (at carbon price \$5 to \$10 per metric tCO₂e). Impacts of recovering primary and secondary forests on the ESs at landscape and local levels are less clear. The results from the global hydrological model, WaterWorld V3.31, employed here showed that the impacts of alternative scenarios on the delivery of watershed services are generally negative at the whole island scale. However, the community surveys showed that local community members strongly believe that declining of watershed services, especially water yield during dry season, is linked to historical events of deforestation and forest degradation.

In terms of local ESs, greatest benefits per household are found where communities are allowed to cultivate and utilize agroforest products (Table 9). Extraction of natural NTFP and timber is important to some, but generally are valued less. Estimation of an aggregate

value of the local ESs in our study area is difficult due to overlapping land use classes and forest functions (Table 3) and also uncertainty of land tenure arrangements. For our analysis, we aggregated the average annual household value of forest ESs for each forest type with the number of households in our study area that have agriculture as their main occupation (Table 11). Our target population for this aggregation was the 23 sub-districts surrounding mount Rinjani. These sub-districts had a population of 1.313 million (with average household size of 3.57) as of 2010 and about 51.5% of population in the area reported agriculture as their main occupation, according to the latest census (BPS/NTB, 2012). The total value of locally provided forest ESs, we aggregate the average household values (Table 9) to the 51.5 % of households (Table 9). The value of local ESs delivered by forests of Lombok is currently estimated at \$16 million to \$18 million annually. Aggregated (undiscounted) over 30 years, the total value ranges from \$486 million to \$564 million.

<Table 11>

To allow a comparison of the carbon values (Table 8) with changes in values of locally provided forest ESs under different land use scenarios, we assume increase in forests in CP and FR scenarios (shown in Table 5 and 6) would be distributed to different forests according to the current ratio.⁷

<Table 12>

Although the predicted changes in locally provided forest ESs values associated with the CP or FR scenarios are approximate, we can demonstrate that these values are higher than the carbon values (\$35.7 - \$69m over 30 years for the Community Partnership scenario and \$38- \$76m for the Forest Restoration scenario).

⁷ Forests in the NTB province includes 20% production forest, 48% protection forest and 32%conservation forest.

Opportunity costs are the forgone economic benefits of alternative land use, in this case dryland agriculture. Communities in the area cultivate various crops, including maize, chili, cassava, peanuts, etc (Collins Higgins Consulting Group, 2012). Lombok is also one of the largest producers of tobacco in Indonesia (Lee et al. 2015). Profitability of dryland agriculture varies a great deal among different varieties of crops and year-to-year. For example, tobacco can go from a net profit to a net loss depending on weather conditions (\$465 to \$1,132/ ha under normal condition to -\$371 to -\$477/ha in a bad year e, g, 2002) (Keyser and Juita, 2005). Net revenue from maize in similar areas has been reported around \$180/ha/yr (Da Silva and Murdolelono. 2010). Table 13 presents opportunity costs of carbon sequestration undiscounted and Net Present Value (NPV) with 10% discount rate over 30-year period per metric tCO₂e with a range of per ha profitability (following the methodology described in White et al. 2010). Opportunity costs are lower than the current carbon price.

<Table 13>

Here we can draw a number of broad conclusions on the ES associations and potential effects of global PES scheme. First, the value of local ESs are potentially greater than that of global ES (carbon) and opportunity costs are low. Thus, carbon PES schemes (such as REDD+) need to be developed in a way to maximize synergies among global and local ESs. Carbon payments can provide the initial capital investment needed for creating nurseries and planting trees, but recovered forests can also provide income overtime for communities to maintain forests. Each community can develop a benefit-sharing mechanism under the partnership agreement (*kemitraan*) with KPHs or through Community Forest arrangement. For example, community D has started tree planting projects with REDD+ demonstration fund facilitated by an NGO (FFI/Indonesia). The species selection was negotiated with the community, and the result was mostly fruit trees planted. Second, higher benefits can be obtained by encouraging secondary forests (retaining artificial gaps in the canopy to 50-60%), while meeting community needs for NTFP, agroforest products and timber. Community partnership scenario is focusing on recovery of secondary forests, which is possible through agroforestry with forest covers. A

previous study in the area shows that carbon stored in agroforestry land with significant forest cover (178 metric ton/ha, Markum et al. 2013), is similar to that in secondary forests (181 metric ton/ha, Table 2). Forest Restoration scenario included additional reforestation to recover primary forests. From the community point of view, primary forest does not generate significant economic revenues, although there may be cultural and religious significance that this study did not capture. Additional carbon payment expected from primary forest can motivate communities to recover primary forests for conservation purposes.

5.2 Data discrepancies: reconciling global modelling and local perceptions

A key debate in ecosystem service assessments relates to identifying what is the most appropriate source of data to measure ecosystem service change (TEEB, 2010). Evaluating watershed services is especially challenging because hydrological impacts can occur anywhere downstream of the site of service production (van Noordwijk et al. 2016). It is not easy to discern the roles of land use change from other influencing factors, such as climate variability, landscape-level changes, and spatial distribution of soil and vegetation types (Bruijnzeel, 2004). In this research, we used both global models (e.g. WaterWorld V3.31) and local knowledge (in-person surveys) to assess the impact of forest management on water regulation. Global models have a wide appeal in that they are usually based on the theoretically sound scientific knowledge and can be applied almost anywhere in the world at relatively low costs. In the absence of long term observation records, collecting local data may require surveys with local stakeholders/communities, which is often based on implicit and experiential knowledge rather than scientific evidence (Christie, 2012). In our research, we found discrepancies between these two data sources, particularly in terms of the predicted impact of forest management on water regulation services.

WaterWorld V3.31 showed that more tree cover decreases baseflow in both dry and wet seasons in most places due to increased evapotranspiration, while increasing baseflow in some places due to enhanced infiltration. This is supported by many studies that indicate higher evapotranspiration of trees than other cover types (Kaimowitz, 2004; Calder, 2001; Van Dijk et al., 2007). The overall effects of both scenarios were negative on watershed

services. However, residents frequently reported contrasting views based on experience and observation. In surveys conducted in Lombok communities in 2002, residents reported that springs had gone dry in response to forest clearing (WWF 2002). According to Pirard (2011), 43% of the large springs surrounding Rinjani dried up in the decade 1992-2002, while approximately 30% of the Mount Rinjani was deforested during the same decade. Klock and Sjah (2012) reported that, during the previous twenty years, more than 400 springs dried up on Mount Rinjani, most likely from deforestation. The Jakarta Post (2014) reported that there are 107 springs currently utilized in Lombok, with many other sources not yet recorded by the government and under the control of local residents. In the above article, a local Village Head is quoted as emphasizing the function of forests as a sponge, absorbing water and releasing it gradually, thus enhancing water regulation and quality. Our community survey also confirm that water regulation was considered important to people living in the forest margins and the follow-up focus group discussions highlighted the strong local belief that retaining and enhancing forest cover protected water supply and water quality.

The prevailing scientific paradigm for linking forests to water has shifted since the early 1980ies when several reviews, both in the temperate zone and the humid tropics, show that there is little empirical evidence for forests storing excess water during wet periods and releasing it during dry periods, so called *sponge theory* (Bosch and Hewlett 1982; Ghimire et al. 2014a; Ilstedt et al. 2016). Since then, many studies supported *trade-off theory*, which means less water yields with increasing tree covers (Ilstedt et al. 2016). Deforestation, especially in the tropics, does contribute soil degradation and increase in impermeable surface, which lead to locally observed negative hydrological effects. However, there is limited evidence for reforestation increasing soil hydraulic conductivities (Ghimire et al. 2013; Ghimire et al. 2014a; Ghimire et al. 2014b). Moreover changes in water resources reflect not only the changes in ecosystem services (modelled here) but also the impacts of farmer behavior of water use and irrigation practices, which was not part of this study. Also, relying only on anecdotal data could lead to an erroneous conclusion regarding changes in spring discharge conditions caused by forest change. As noted above, illegal logging, encroachment and occupation reached its peak after the fall of

the Suharto regime in 1998. Loss of forest cover notwithstanding, climate variation could have had a bearing on residents' perception of the effects of forest clearing. Long-term precipitation records shows that there are a great deal variations in precipitation during dry season among different locations and also years leading up to 1998 were dry, especially around the Mataram city in low elevation. Figure 6 shows average precipitation records from six weather stations around the city of Mataram and four weather stations near the survey locations around Rinjani Mountain. It is very possible that declining spring discharge was more directly related to climate than to land use change. Furthermore, the existence of the PES mechanism between the city of Mataram and the communities in their upper watershed area may have raised expectation of forest-margin communities that they may be able to be compensated for managing forest for watershed services that they provide. It may be especially true for the community D that gained Community Forest recognition and their forest represents important watershed for another city (city of Praya).

<Figure 6>

WaterWorld V3.31 simulates the impacts of forests versus other land uses on hydrological impacts based on high resolution remotely sensed data. It is a very detailed process model developed specifically for data poor mountainous and tropical environments. However, its results can not be field-validated without long-term spring discharge measurement data, and are not without limitations. Change in land cover and forest canopy structure have complex effects on fog input, rainfall interception, throughfall, stemflow, infiltration and runoff generation (Bruijnzeel et al., 2011; Dietz et al., 2006, Bruijnzeel et al., 2004). Some have argued that in contrast to other land use cover types, natural and recovered tropical rainforests throughout the world exhibit greater leaf litter, soil organic matter, and soil bioturbation by roots and fauna, as well as less soil surface sealing due to rainsplash, soil compaction by farm equipment, and impervious surface as part of infrastructure, all of which allow for enhanced infiltration and reduced soil erosion (Kumagai et al. 2009, Hairah et al. 2006, Bruijnzeel et al. 2004, Calder 2001, Mapa 1995). The net result of enhanced infiltration beneath recovered forest can be greater groundwater recharge, which can lead

to improved dry season baseflow (Dias et al. 2015, Ogden et al. 2013, Peña-Arancibia et al. 2012, Bruijnzeel et al. 2006, Bruijnzeel et al. 2004). Forests do tend to increase evapotranspiration substantially compared with rain-fed agriculture and even higher infiltration rates cannot compensate for less water being available for infiltration and runoff. However, this basic assumption may be problematic in a tropical setting where atmospheric moisture is abundant; low vapor pressure deficit may result in reforestation having a negligible effect on evapotranspiration (Brauman 2012). Malder et al. (2013) argued that the data to formulate hydrological effects of land use change in global models are often generated outside the tropics with stable soil conditions and there is “complete lack of research on how forestation on degraded land affect hydrological functioning at the landscape scale.” Empirical long-term spring discharge measurement data are needed to compliment and refine global models based on globally available datasets, in order to accurately evaluate land management practices that enhance watershed services (Wohl et al. 2012, Jose 2009, Locatelli and Vignola 2009).

What is clear from the above discussions is that there are number of factors that might affect the accuracy of both the global models and local opinions. Simply focusing on increasing tree covers can have negative impacts on watershed services and set up false expectations among local communities. For example, empirical studies in other seasonal dry tropics showed that reforestation with pine species in densely populated areas did little to increase soil hydraulic conductivities while increasing water uses of vegetation (compared to pasture) (Ghimire et al. 2013; Ghimire et al. 2014a; Ghimire et al. 2014b). Another study showed that hydrological benefits of reforestation can be maximized by considering the rates of evapotranspiration of different tree species, as well as tree size, age and density in planning reforestation projects (Ilstedt et al. 2016). Thus, global PES schemes must consider further details within a land use class (e.g. species selection, tree density, soil management, and landscape configurations) and measures to mitigate potential negative impacts.

5.3 Tradeoffs and synergies between global and local ecosystem services

Globally, simply ending the land use, passive restoration, has been shown to be more cost-effective than active restoration (Meli et al. 2017). However, in a densely populated region with complex social dynamics, protection of forest as carbon stock would be costly and ineffective (Skutsch et al. 2011). In both land use change scenarios, there is potential for developing forest carbon projects in the study area. Although on-site opportunity costs were low, social and indirect costs can be substantial (White et al., 2010). Most of the global forest carbon projects are financed as input-based projects, which often set a flat-rate payment per hectare under a contractual agreement of inputs to increase carbon stock (e.g., not cutting trees, tree planting or other management activities) (Wunder, 2008; Skutsch et al., 2011). Input-based carbon projects allow the inputs (e.g. agreed management actions) to be negotiated between project proponents and local communities, which makes the projects less politically contentious and allows broader management goals to be addressed (Skutsch et al., 2011). However, input-based projects would likely generate fewer carbon credits overall while making it difficult to trace carbon to project activities (Skutsch et al., 2011). Lack of reporting on actual performance of existing projects, in terms of carbon sequestration, poses a serious problem for the future of global carbon financing (Fischer et al., 2016).

We previously advocated for an input-based mechanism with readiness activities for capacity building of both institutions and communities in the study area (Kim et al. 2016). The results of this study show that simply increasing tree cover is not enough for enhancing ES at all scales. Reforestation to increase carbon stock without considering the landscape as a whole can have negative impacts on watershed services (e.g. reduced runoff, and concentrated pollutants downstream from the remaining agricultural lands). In addition, implementing reforestation projects without consideration for local livelihoods can be detrimental to forest-margin communities. Thus the details of agreed-upon management actions would dictate the nature of association among different ESs.

Previous studies argued that global forest carbon projects are unlikely to succeed without addressing food, energy and water provisions at the local level (Minang and van Noordwijk,

2013; van Noordwijk et al., 2016). Indeed, the findings from our community study demonstrate that local people obtain a wide range of benefits from forests. Mixed agroforestry systems can be a key strategy for increasing the multi-functionality of land uses (Minang et al., 2014) as well as enhancing the diversity of local communities' livelihood options (Hoang et al., 2014). Potential values of agroforestry systems for integrating forests into a multifunctional landscape have been recognized, although the benefits may vary depending on practices and landscape configurations (Table 9; Dewi et al., 2013; Prabhu et al. 2015). It has been shown that intermediate tree cover can maximize ground water recharge in seasonally dry tropics (*optimal tree cover theory*)(Ilstedt et al. 2016). Impacts of agroforestry systems on the landscape's ability to provide watershed services vary depending on species selection of crops and shade trees and different cultivation practices employed (Condon et al. 2002; Thierfelder et al. 2009). For example, different tropical tree species have shown a wide range of production rates per cost of water loss by transpiration (Cernusak et al. 2007) and different root depths for promoting soil infiltration of rainfall (Ghestem et al. 2011). Local communities that we surveyed also recognized specific "watershed trees" e.g. Beringin (*Ficus benjamina*), where soils underneath were observed to be more moist, compared to other fast growing species, e.g. Sengon (*Albizia chinensis*). Also the amount of water needed to produce different agroforestry crops varies greatly. For example, coffee and cacao tend to have high water footprint (about 22,900 m³/ton for coffee and 9,414 m³/ton for cacao), compared to other crops (e.g. 514 m³/ton for cassava) (Bulsink et al. 2009). Thus it is essential for forest carbon projects to consider the effects of increasing tree covers, along with species, size, and age distribution, on a range of ESs in the landscape and mitigate potential negative impacts. van Noordwijk et al. (2016) discussed several metrics for developing mitigation actions through agroforestry that can enhance different watershed services, including water yield, water flow and water quality, while improving local livelihoods. The plausible actions that can be incorporated into forest carbon projects include replacing fast growing tree plantations with low-evapotranspiration species and increasing presence of deep rooted trees while promoting litter layers and agricultural practices that increase infiltration and soil water content, enhancing sediment filter strips in fields and across

landscape matrix, as well as protecting river banks, riparian zones and landslide-prone slopes, springs and sources of domestic water use.

It is clear from the community surveys that the value of forest ESs to local communities is significant but vary by locations. Although it is difficult to fully untangle the underlying reasons for this, these differences are reflective of different designated functions of forest, suitability of land for agroforestry, and the security of land tenure. Community partnership scenario focused on recovery of secondary forests through agroforestry to provide food, energy and livelihood options for local communities. However, the synergy among global, landscape and local ESs can be created only if the clear accountability can be established for maintaining the threshold of forest covers (for carbon accounting) with specific species selection and agroforestry practices to increase soil infiltration and water use efficiency (for watershed services). Although the Forest Restoration scenario adds recovery of primary forests, local communities may lack motivation for restoration activities for ecological benefits alone. Global PES, such as REDD+, can help establishing technical guidelines for agroforestry practices that maximize carbon and watershed benefits, as well as developing community monitoring schemes, while promoting ecological restoration of primary forest with added carbon values under Forest Restoration scenario.

6 Conclusions

In this paper, we assessed realistic forest management scenarios for reforestation in Eastern Indonesia and their effects on both global and local ES provisions. We have demonstrated that reforestation to increase carbon, i.e. global, ex-situ, ecosystem services, can have varying impacts on those ESs recognized locally. In particular, our results point to the significance of water regulation, agroforest products, and non-timber forest products to local communities. To create a sustainable local solution, we need to go beyond the zero-sum argument of livelihoods versus conservation. We demonstrated how global PES, such as REDD+, and landscape level PES, such as payment for watershed services, can help create, not dictate, such solution through agroforestry that meets global, landscape and local demands for ESs.

1457
1458
1459 679
1460

1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512

7 References

- Bonan, G.B. 2008. Forests and climate change: forcings, feedbacks, and the climate benefits of forests. *Science*, 320, 1444–1449.
- Bae, J.S., Kim, C., Kim, Y.-S., Latifah, S., Afifi, M., Fisher, L.A., Lee, S.M., Kim, I.-A., Kang, J., Kim, R., Kim, J.S., 2014. Opportunities for implementing REDD+ to enhance sustainable forest management and improve livelihoods in Lombok, NTB, Indonesia. Working Paper 151. CIFOR. Bogor, Indonesia
- Biermann, F., 2010. Beyond the intergovernmental regime: recent trends in global carbon governance. *Current Opinion in Environmental Sustainability*, 2(4), pp.284-288.
- Blom, B., Sunderland, T., Murdiyarso, D., 2011. Getting REDD to work locally, lessons learned from integrated conservation and development projects. *Environ. Sci. Policy* 13, 164–172.
- Bosch, J.M. and Hewlett, J.D., 1982. A review of catchment experiments to determine the effect of vegetation changes on water yield and evapotranspiration. *Journal of hydrology*, 55(1-4), pp.3-23.
- Brauman, K.A., Freyberg, D.L., and Daily, G.C. 2012. Potential evapotranspiration from forest and pasture in the tropics: A case study in Kona, Hawai'i. *Journal of Hydrology*, Volumes 440–441: 52–61.
- Bruijnzeel, L.A., M. Mulligan, and F.S. Scantena. 2011. Hydrometeorology of tropical montane cloud forests: emerging patterns. *Hydrological Processes*, 25, 465-498.
- Bruijnzeel, L.A., Burkard, R., Carvajal, A., Frumau, K.F.A., Köhler, L., Mulligan, M., Schellekens, J., Schmid, S. and Tobón-Marin, C., 2006. Hydrological impacts of converting tropical montane cloud forest to pasture, with initial reference to northern Costa Rica. DFID Project Report
- Bruijnzeel, L.A. 2004. Hydrological functions of tropical forests: not seeing the soil for the trees? *Agriculture, Ecosystems and Environment* 104 (2004) 185–228.
- BPS, 2016. . Indonesia Human Development Index. Available at: http://www.bps.go.id/eng/tab_sub/view.php?tabel=1&daftar=1&id_subyek=26&no_tab=2. (accessed Aug. 10 2016).
- BPS-NTB, 2015. Nusa Tenggara Barat dalam Angka (NTB Statistics) Tahun 2014. Badan Pusat Statistik Provinsi Nusa Tenggara Barat. Mataram.
- BPS, 2012. Regional profile, Nusa Tenggara Barat, Available online: <http://www.bps.go.id/profile/ntb.shtml> (accessed Aug. 10 2013).

- Bulsink, F., Hoekstra, A.Y. and Booij, M.J., 2009. The water footprint of Indonesian provinces related to the consumption of crop products, Value of Water Research Report Series No. 37.
- Bruckner, M., Fischer, G., Tramberend, S. and Giljum, S., 2015. Measuring telecouplings in the global land system: a review and comparative evaluation of land footprint accounting methods. *Ecological Economics*, 114, pp.11-21.
- Calder, I.R. 2001. Canopy processes: implications for transpiration, interception and splash induced erosion, ultimately for forest management and water resources. *Plant Ecology*, 153: 203–214, 2001.
- Cash, D., Adger, W.N., Berkes, F., Garden, P., Lebel, L., Olsson, P., Pritchard, L. and Young, O., 2006. Scale and cross-scale dynamics: governance and information in a multilevel world. *Ecology and society*, 11(2).
- Cernusak, L.A., Aranda, J., Marshall, J.D. and Winter, K., 2007. Large variation in whole-plant water-use efficiency among tropical tree species. *New Phytologist*, 173(2), pp.294-305.
- Christie M, Fazey I, Cooper R, Hyde H and Kenter JO. 2012. An Evaluation of Monetary and Non-monetary Techniques for Assessing the Importance of Biodiversity and Ecosystem Services to People in countries with developing economies. *Ecological Economics*, 83, 69-80.
- Collins Higgins Consulting Group. 2012. Project Analysis of Agribusiness Development Opportunities in Easter Indonesia – Socio-Economic Review. Prepared for the Australian Centre for International Agricultural Research by Collins Higgins Consulting Group Pty Ltd.
- Condon, A.G., Richards, R.A., Rebetzke, G.J. and Farquhar, G.D., 2002. Improving intrinsic water-use efficiency and crop yield. *Crop Science*, 42(1), pp.122-131.
- Da Silva, H. and Murdolelono, B. 2010. Feasibility of OPV Srikandi Maize for Overcoming Productivity and Food Security Problems in East Nusa Tenggara, Indonesia. In 'Maize for Asia: emerging trends and technologies. Proceeding of the 10th Asian Regional Maize Workshop, Makassar, Indonesia, 20-23 October 2008', ed. by P.H. Zaidi, M. Azrai and K.V. Pixley. CIMMYT
- Daily, G.C., Polasky, S., Goldstein, J., Kareiva, P.M., Mooney, H.A., Pejchar, L., Ricketts, T.H., Salzman, J. and Shallenberger, R., 2009. Ecosystem services in decision making: time to deliver. *Frontiers in Ecology and the Environment*, 7(1), pp.21-28.
- de Groot, R.S., Alkemade, R., Braat, L., Hein, L. and Willemen, L., 2010. Challenges in integrating the concept of ecosystem services and values in landscape planning, management and decision making. *Ecological complexity*, 7(3), pp.260-272.

- Dewi, S., van Noordwijk, M., Ekadinata, A. and Pfund, J.L., 2013. Protected areas within multifunctional landscapes: Squeezing out intermediate land use intensities in the tropics?. *Land Use Policy*, 30(1), pp.38-56.
- Dias, L.C.P., Macedo, M.N., Costa, M.H., Coe, M.T. and Neill, C., 2015. Effects of land cover change on evapotranspiration and streamflow of small catchments in the Upper Xingu River Basin, Central Brazil. *Journal of Hydrology: Regional Studies*, 4, pp.108-122.
- Dietz, J., Hölscher, D. and Leuschner, C., 2006. Rainfall partitioning in relation to forest structure in differently managed montane forest stands in Central Sulawesi, Indonesia. *Forest Ecology and Management*, 237(1), pp.170-178.
- Diswandi, D., 2017. A hybrid Coasean and Pigouvian approach to Payment for Ecosystem Services Program in West Lombok: Does it contribute to poverty alleviation?. *Ecosystem Services*, 23, pp.138-145.
- Enrici, A. and Hubacek, K., 2016. Business as usual in Indonesia: governance factors effecting the acceleration of the deforestation rate after the introduction of REDD+. *Energy, Ecology and Environment*, 1(4), pp.183-196.
- Food and Agricultural Organization(FAO), 2010. Global Forest Resources Assessment 2010, Main report. FAO Forestry Paper No. 163. Rome, FAO, 340-41.
- Fazey, I, Kesby, M, Evely, A, Latham, I, Wagatora, D, Hagasua, J-E, Reed, M, S, Christie, M 2010. A three-tiered approach to participatory vulnerability assessment in the Solomon Islands. *Global Environmental Change*. 20, 713-728
- Fischer, R., Hargita, Y. and Günter, S., 2016. Insights from the ground level? A content analysis review of multi-national REDD+ studies since 2010. *Forest Policy and Economics*, 66, pp.47-58.
- Geist, H.J. and Lambin, E.F., 2002. Proximate causes and underlying driving forces of tropical deforestation: Tropical forests are disappearing as the result of many pressures, both local and regional, acting in various combinations in different geographical locations. *BioScience*, 52(2), pp.143-150.
- Ghestem, M., Sidle, R.C. and Stokes, A., 2011. The influence of plant root systems on subsurface flow: implications for slope stability. *BioScience*, 61(11), pp.869-879.
- Ghimire, C.P., Bonell, M., Bruijnzeel, L.A., Coles, N.A., Lubczynski, M.W., 2013. Reforesting severely degraded grassland in the Lesser Himalaya of Nepal: effects on soil hydraulic conductivity and overland flow production. *Journal of Geophysics Research- Earth Surface*. 118, <http://dx.doi.org/10.1002/2013JF002888>.

- Ghimire, C.P., Bruijnzeel, L.A., Lubczynski, M.W. and Bonell, M., 2014a. Negative trade-off between changes in vegetation water use and infiltration recovery after reforestation degraded pasture land in the Nepalese Lesser Himalaya. *Hydrology and Earth System Sciences*, 18(12), p.4933
- Ghimire, C.P., Bruijnzeel, L.A., Bonell, M., Coles, N., Lubczynski, M.W. and Gilmour, D.A., 2014b. The effects of sustained forest use on hillslope soil hydraulic conductivity in the Middle Mountains of Central Nepal. *Ecohydrology*, 7(2), pp.478-495.
- Goetz, S.J., Hansen, M., Houghton, R.A., Walker, W., Laporte, N. and Busch, J., 2015. Measurement and monitoring needs, capabilities and potential for addressing reduced emissions from deforestation and forest degradation under REDD+. *Environmental Research Letters*, 10(12), p.123001.
- Goldstein, A., Neyland, E. and Bodnar, E., 2015. Converging at the crossroads State of forest carbon finance 2015. *Forest Trends' Ecosystem Marketplace*. Washington, DC.
- Gómez-Baggethun E, de Groot R, Lomas PL, Montes C. 2010. The history of ecosystem services in economic theory and practice: from early notions to markets and payment schemes. *Ecological Economics* 69(6): 1209–1218. DOI: 10.1016/j.ecolecon.2009.11.007
- González-Esquivel, C.E., Gavito, M.E., Astier, M., Cadena-Salgado, M., del-Val, E., Villamil-Echeverri, L., Merlín-Urbe, Y. and Balvanera, P., 2015. Ecosystem service trade-offs, perceived drivers, and sustainability in contrasting agroecosystems in central Mexico. *Ecology and Society*, 20(1).
- Gleeson, T., Smith, L., Moosdorf, N., Hartmann, J., Dürr, H.H., Manning, A.H., van Beek, L.P. and Jellinek, A.M., 2011. Mapping permeability over the surface of the Earth. *Geophysical Research Letters*, 38(2).
- Grossman, J., 2015. Ecosystem service trade-offs and land use among smallholder farmers in eastern Paraguay. *Ecology and Society*, 20(1).
- Haines-Young, R., Potschin, M. and Kienast, F., 2012. Indicators of ecosystem service potential at European scales: mapping marginal changes and trade-offs. *Ecological Indicators*, 21, pp.39-53.
- Hansen, M.C., Potapov, P.V., Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A., Thau, D., Stehman, S.V., Goetz, S.J., Loveland, T.R. and Kommareddy, A., 2013. High-resolution global maps of 21st-century forest cover change. *science*, 342(6160), pp.850-853.
- Hamrick, K. and Goldstein, A., 2016. Raising ambition: State of the voluntary carbon markets 2016. *Forest Trends' Ecosystem Marketplace*. Washington, DC.

- Hairiah, K., Sulistyani, H., Suprayogo, D., Purnomosidhi, P., Widodo, R.H. and Van Noordwijk, M., 2006. Litter layer residence time in forest and coffee agroforestry systems in Sumberjaya, West Lampung. *Forest Ecology and Management*, 224(1), pp.45-57.
- Hein, L., Van Koppen, K., De Groot, R.S. and Van Ierland, E.C., 2006. Spatial scales, stakeholders and the valuation of ecosystem services. *Ecological economics*, 57(2), pp.209-228.
- Hoang, M.H., Namirembe, S., van Noordwijk, M., Catacutan, D., Öborn, I., Perez-Teran, A.S., Nguyen, H.Q. and Dumas-Johansen, M.K., 2014. Farmer portfolios, strategic diversity management and climate-change adaptation–implications for policy in Vietnam and Kenya. *Climate and Development*, 6(3), pp.216-225.
- Ilstedt, U., Bargués Tobella, A., Bazié, H.R., Bayala, J., Verbeeten, E., Nyberg, G., Sanou, J., Benegas, L., Murdiyarso, D., Laudon, H., Sheil, D., Malmer, A., 2016. Intermediate tree cover can maximize groundwater recharge in the seasonally dry tropics. *Sci. Rep.* 6, 21930.
- Intarini, D.Y., Resosudarmo, I.A.P., Komalasari, M., Ekaputri, A.E., 2015. Ketapang Community Carbon Pools, Ketapang, West Kalimantan, Indonesia, in: Sills E.O., tmdja, S., de Sassi, C., Duchelle, A.E., Kweka, D., Resosudarmo, I.A.P., Sunderlin W.D. (Eds.), REDD+ on the Ground: A Case Book of Subnational Initiatives around the World, CIFOR. Bogor, Indonesia
- Jakarata Post. 2014. Preserving Lombok's natural water sources. The Jakarta Post, Tuesday, June 24, 2014, 12:24 pm.
- Jang, S.K., Bae, J. S., 2014. New Forest Governance in Indonesia: A Forest Partnership between Forest Management Units and Local Communities. *Southeast Asia Study* 24, 95-144
- Jose, S. 2009. Agroforestry for ecosystem services and environmental benefits: an overview. *Agroforestry Systems*, 76 (1) 1-10.
- Kaimowitz, D., 2004. Useful myths and intractable truths: the politics of the link between forests and water in Central America. *Forest, Water and People in the Humid Tropics: Past, Present and Future Hydrological Research for Integrated Land and Water Management*, pp.86-98.
- Keyser, J.C. and Ratna Juita, N., 2005. Smallholder tobacco growing in Indonesia: Costs and profitability compared with other agricultural enterprises. *The World Bank*
- Kim, Y.S., Bae, J.S., Fisher, L.A., Latifah, S., Afifi, M., Lee, S.M. and Kim, I.A., 2016. Indonesia's forest management units: effective intermediaries in REDD+ implementation?. *Forest Policy and Economics*, 62, pp.69-77.

- Klock, J., and Sjah, T. 2011. Farmer water management strategies for dry season water shortages in central Lombok, Indonesia. *Natural Resources*, 2011, 2, 114-124. doi:10.4236/nr.2011.
- Kumagai, T.O., Yoshifuji, N., Tanaka, N., Suzuki, M. and Kume, T., 2009. Comparison of soil moisture dynamics between a tropical rain forest and a tropical seasonal forest in Southeast Asia: Impact of seasonal and year-to-year variations in rainfall. *Water Resources Research*, 45(4).
- Lambin, E.F. and Meyfroidt, P., 2011. Global land use change, economic globalization, and the looming land scarcity. *Proceedings of the National Academy of Sciences*, 108(9), pp.3465-3472.
- Lee, S.M., Kim, Y.S., Jaung, W., Latifah, S., Afifi, M. and Fisher, L.A., 2015. Forests, fuelwood and livelihoods—energy transition patterns in eastern Indonesia. *Energy Policy*, 85, pp.61-70.
- Leimona, B., Lusiana, B., van Noordwijk, M., Mulyoutami, E., Ekadinata, A. and Amaruzaman, S., 2015. Boundary work: Knowledge co-production for negotiating payment for watershed services in Indonesia. *Ecosystem Services*, 15, pp.45-62.
- Liu, J., Mooney, H., Hull, V., Davis, S.J., Gaskell, J., Hertel, T., Lubchenco, J., Seto, K.C., Gleick, P., Kremen, C. and Li, S., 2015. Systems integration for global sustainability. *Science*, 347(6225), p.1258832..
- Locatelli, B. and Vignola, R., 2009. Managing watershed services of tropical forests and plantations: can meta-analyses help?. *Forest Ecology and Management*, 258(9), pp.1864-1870.
- Maes, J., Paracchini, M.L., Zulian, G., Dunbar, M.B. and Alkemade, R., 2012. Synergies and trade-offs between ecosystem service supply, biodiversity, and habitat conservation status in Europe. *Biological conservation*, 155, pp.1-12.
- Malmer, A., Murdiyarso, D. Bruijnzeel, L.A. and Ilstedt, U. 2010. Carbon sequestration in tropical forests and water: a critical look at the basis for commonly used generalizations. *Global Change Biology*, 16(2), pp.599-604.
- Mapa, R.B., 1995. Effect of reforestation using *Tectona grandis* on infiltration and soil water retention. *Forest Ecology and Management*, 77(1-3), pp.119-125.
- Margono, B.A., Potapov, P.V., Turubanova, S., Stolle, F. and Hansen, M.C., 2014. Primary forest cover loss in Indonesia over 2000-2012. *Nature Climate Change*, 4(8), pp.730-735.
- Markum, M., Soesilaningsih, E.A., Suprayogo, D. and Hairiah, K., 2013 Contribution of Agroforestry System in Maintaining carbon stocks and reducing emission rate at

1849
1850
1851 888 Jangkok Watershed, Lombok island. AGRIVITA, Journal of Agricultural Science,
1852 889 35(3), pp.207-217.
1853
1854
1855 890 Mastrangelo, M.E. and Laterra, P., 2015. From biophysical to social-ecological trade-offs:
1856 891 integrating biodiversity conservation and agricultural production in the Argentine
1857 892 Dry Chaco. Ecology and Society, 20(1).
1858
1859 893 MA (Millennium Ecosystem Assessment). 2005. Ecosystems and Human Well-being:
1860 894 Synthesis. Island Press, Washington, DC.
1861
1862 895 Meli, P., Holl, K.D., Benayas, J.M.R., Jones, H.P., Jones, P.C., Montoya, D. and Mateos, D.M.,
1863 896 2017. A global review of past land use, climate, and active vs. passive restoration
1864 897 effects on forest recovery. PloS one, 12(2), p.e0171368.
1865
1866 898 Meyfroidt, P., Lambin, E.F., Erb, K.H. and Hertel, T.W., 2013. Globalization of land use:
1867 899 distant drivers of land change and geographic displacement of land use. Current
1868 900 Opinion in Environmental Sustainability, 5(5), pp.438-444.
1870
1871 901 Minang, P.A., van Noordwijk, M., Freeman, O.E., Mbow, C., de Leeuw, J. and Catacutan, D.
1872 902 eds., 2014. Climate-smart landscapes: multifunctionality in practice. ASB
1873 903 Partnership for The Tropical Forest margins.
1874
1875 904 Minang, P.A. and van Noordwijk, M., 2013. Design challenges for achieving reduced
1876 905 emissions from deforestation and forest degradation through conservation:
1877 906 leveraging multiple paradigms at the tropical forest margins. Land Use Policy, 31,
1878 907 pp.61-70.
1879
1880 908 Motel, P.C., Pirard, R. and Combes, J.L., 2009. A methodology to estimate impacts of
1881 909 domestic policies on deforestation: Compensated Successful Efforts for "avoided
1882 910 deforestation"(REDD). Ecological Economics, 68(3), pp.680-691.
1883 910
1884
1885 911 Mora, F., Balvanera, P., García-Frapolli, E., Castillo, A., Trilleras, J., Cohen-Salgado, D. and
1886 912 Salmerón, O., 2016. Trade-offs between ecosystem services and alternative
1887 913 pathways toward sustainability in a tropical dry forest region. Ecology and Society,
1888 914 21(4).
1889
1890 915 Mouchet, M.A., Lamarque, P., Martín-López, B., Crouzat, E., Gos, P., Byczek, C. and Lavorel, S.,
1891 916 2014. An interdisciplinary methodological guide for quantifying associations
1892 917 between ecosystem services. Global Environmental Change, 28, pp.298-308.
1893 917
1894
1895 918 Mulligan, M. 2013. WaterWorld: a self-parameterising, physically based model for
1896 919 application in data-poor but problem-rich environments globally. IWA Publishing,
1897 920 Hydrology Research, 44.5: 748-769.
1898
1899 921 Mulligan, M. 2010. Modeling the tropics-wide extent and distribution of cloud forest and
1900 922 cloud forest loss, with implications for conservation priority in Bruijnzeel, L.A.,
1901
1902
1903
1904

1905
1906
1907 923 Scatena, F.N., Hamilton, L.S. (editors), Tropical Montane Cloud Forests: Science for
1908 924 Conservation and Management. Cambridge University Press, Cambridge. pp14-39
1909
1910
1911 925 Ogden, F. L., Crouch, T.D., Stallard, R.F., and Hall, J.S. 2013. Effect of land cover and use on
1912 926 dry season river runoff, runoff efficiency, and peak storm runoff in the seasonal
1913 927 tropics of Central Panama. *Water Resources Research*, 49 (12) 8443–8462.
1914
1915 928 Pan Y., R.A. Birdsey, J. Fang, R. Houghton, P.E. Kauppi, W.A. Kurz, O.L. Phillips, A. Shvidenko,
1916 929 S.L. Lewis, J.G. Canadell, P. Ciais, R.B. Jackson, S.W. Pacala, A.D. McGuire, S. Piao, A.
1917 930 Rautiainen, S. Sitch, D. Hayes. 2011. A large and persistent carbon sink in the world's
1918 931 forests. *Science* 333: 988-993.
1919
1920 932 Peña-Arancibia, J.L., van Dijk, A.I.J., M., Guerschman, J.P. Mulligan, M. Bruijnzeel, A.L. and
1921 933 McVicar, T.R. 2012. Detecting changes in streamflow after partial woodland clearing
1922 934 in two large catchments in the seasonal tropics. *Journal of Hydrology* 416–417: 60–
1923 935 71.
1924
1925
1926 936 Pirard, R. 2012. Payments for Environmental Services (PES) in the public policy landscape:
1927 937 “Mandatory” spices in the Indonesian recipe. *Forest Policy and Economics*. 18:23-29
1928
1929 938 Pistorius T. 2012. From RED to REDD+: The evolution of a forest-based mitigation
1930 939 approach for developing countries. *Current Opinion in Environmental Sustainability*
1931 940 4(6):638-45.
1932
1933 941 Prabhu, R., Barrios, E., Bayala, J., Diby, L., Donovan, J., Gyau, A., Gaudal, L., Jamnadass, R.,
1934 942 Kahia, J., Kehlenbeck, K. and Kindt, R., 2015. Agroforestry: Realising the promise of
1935 943 an agroecological approach. In *Agroecology for Food Security and Nutrition:*
1936 944 *Proceedings of the FAO International Symposium, Rome, FAO* (pp. 201-224).
1937
1938
1939 945 Prasetyo, F.A., Aritta, S., Purwanto, Hakim, H., 2009. Making Policies Work for Payment for
1940 946 Environmental Services (PES), Evaluation of the Experience of Formulating
1941 947 Conservation Policies in Districts of Indonesia. *Journal of Sustainable Forestry*. 28,
1942 948 415–433
1943
1944 949 Republic of Indonesia (RoI). 2007. Government Regulation No. 6/2007 Forest Arrangement
1945 950 and Establishment of Forest Management Plan as well as Forest Exploitation (in
1946 951 Indonesian). Republic of Indonesia, Jakarta. Indonesia
1947
1948
1949 952 Resosudarmo, I.A.P., 2004. Closer to people and trees: will decentralisation work for the
1950 953 people and the forests of Indonesia?. *The European Journal of Development*
1951 954 *Research*, 16(1), pp.110-132.
1952
1953 955 Rosenbarger, A., Gingold, B., Prasodjo, R., Alisjahbana, A., Putraditama, A. and Tresya, D.,
1954 956 2013. How to change legal land use classifications to support more sustainable palm
1955 957 oil in Indonesia. World Resources Institute, Washington, DC.
1956
1957
1958
1959
1960

1961
1962
1963
1964 958 RoI. 1999. Law No. 41 of 1999 on Forestry. Republic of Indonesia, Jakarta. Indonesia
1965
1966 959 Rodríguez, J., Beard Jr, T.D., Bennett, E., Cumming, G., Cork, S., Agard, J., Dobson, A. and
1967 960 Peterson, G., 2006. Trade-offs across space, time, and ecosystem services. *Ecology*
1968 961 and society, 11(1).
1969
1970 962 Sahide, M. A. K., S. Supratman, A. Maryudi, Y.-S. Kim, and L. Giessen. 2016. Decentralisation
1971 963 Policy as Recentralisation Strategy: Forest Management Units and Community
1972 964 Forestry in Indonesia. *International Forestry Review*, 18(1), 78-95.
1973
1974 965 Schweizer, R., Dupuis, J. and De Buren, G., 2016. Environmental innovation strategies:
1975 966 When and why NGOs go beyond public regulations. *Environmental Politics*, 25(5),
1976 967 pp.899-920.
1977
1978 968 Seto, K.C., Reenberg, A., Boone, C.G., Fragkias, M., Haase, D., Langanke, T., Marcotullio, P.,
1979 969 Munroe, D.K., Olah, B. and Simon, D., 2012. Urban land teleconnections and
1980 970 sustainability. *Proceedings of the National Academy of Sciences*, 109(20), pp.7687-
1982 971 7692.
1983
1984 972 Skutsch, M., Vickers, B., Georgiadou, Y. and McCall, M., 2011. Alternative models for carbon
1985 973 payments to communities under REDD+: A comparison using the Polis model of
1986 974 actor inducements. *Environmental Science & Policy*, 14(2), pp.140-151.
1987
1988 975 Simula, M. 2010. Analysis of REDD+ financing gaps and overlaps. REDD+ partnership.
1989 976 <http://reddpluspartnership.org/65524/en/> (accessed 14 November 2011).
1990
1991 977 Stevens C, Winterbottom R, Springer J and Reyntar K. 2014. Securing Rights, Combating
1992 978 Climate Change: How Strengthening Community Forest Rights Mitigates Climate
1993 979 Change. Washington DC: World Resources Institute and Rights and Resources
1995 980 Initiative.
1996
1997 981 TEEB, 2010. The Economics of Ecosystems and Biodiversity: Ecological and Economic
1998 982 Foundations. Edited by Pushpam Kumar Earthscan, London and Washington.
1999
2000 983 Thierfelder, C. and Wall, P.C., 2009. Effects of conservation agriculture techniques on
2001 984 infiltration and soil water content in Zambia and Zimbabwe. *Soil and tillage*
2002 985 *research*, 105(2), pp.217-227.
2003
2004 986 Turner, B.L., Lambin, E.F. and Reenberg, A., 2007. The emergence of land change science for
2005 987 global environmental change and sustainability. *Proceedings of the National*
2006 988 *Academy of Sciences*, 104(52), pp.20666-20671.
2007
2008
2009 989 Turner, K.G., Odgaard, M.V., Bøcher, P.K., Dalgaard, T. and Svenning, J.C., 2014. Bundling
2010 990 ecosystem services in Denmark: Trade-offs and synergies in a cultural landscape.
2011 991 *Landscape and Urban Planning*, 125, pp.89-104.
2012
2013
2014
2015
2016

2017
2018
2019
2020 992 UN-REDD Programme, 2017, About REDD+. Geneva, UN-REDD Programme. Available
2021 993 online: <http://www.un-redd.org/> (accessed March. 10 2017).
2022
2023 994 van Dijk, A.I. and Keenan, R.J., 2007. Planted forests and water in perspective.
2024
2025 995 van Noorwijk, M., Y.-S. Kim, B. Leimona, K. Hairiah, L.A. Fisher. 2016. Metrics of Water
2026 996 Security, Adaptive Capacity, and Agroforestry in Indonesia. *Current Opinion in*
2027 997 *Environmental Sustainability* 21: 1-8 DOI: 10.1016/j.cosust.2016.10.00
2028
2029 998 VCS. 2012 (updated in 2013) Estimation of baseline carbon stock changes and greenhouse
2030 999 gas emissions from unplanned deforestation (BL-UP), v3.2 Available [http://v-c-](http://v-c-s.org/methodologies/VM0007)
2031 1000 [s.org/methodologies/VM0007](http://v-c-s.org/methodologies/VM0007) Last accessed Nov. 2013.
2032
2033 1001 White, D., Minang, P., Agus, F., Borner, J., Hairiah, K., Gockowski, J., Hyman, G., Robiglia, V.,
2034 1002 Swallow, B., Velarde, S. and van Noordwijk, M., 2010. Estimating the opportunity
2035 1003 costs of REDD+, a training manual. World Bank, Washington (D.C.), USA.
2036
2037
2038 1004 Willemsen, L., Hein, L., van Mensvoort, M.E. and Verburg, P.H., 2010. Space for people, plants,
2039 1005 and livestock? Quantifying interactions among multiple landscape functions in a
2040 1006 Dutch rural region. *Ecological Indicators*, 10(1), pp.62-73.
2041
2042 1007 Wohl, E., Barros, A., Brunsell, N., Chappell, N.A., Coe, M., Giambelluca, T., Goldsmith, S.,
2043 1008 Harmon, R., Hendrickx, J.M., Juvik, J. and McDonnell, J., 2012. The hydrology of the
2044 1009 humid tropics. *Nature Climate Change*, 2(9), pp.655-662.
2045
2046 1010 Wunder, S., Engel, S., Pagiola, S., 2008. Taking stock: A comparative analysis of payments for
2047 1011 environmental services programs in developed and developing countries. *Ecological*
2048 1012 *Economics*. 65, 834-852.
2049
2050
2051 1013 WWF. 2008. Studi Analisis Hidrologis Dan Perubahan Tutupan Lahan (land use land cover
2052 1014 change) Kawasan Gunung Rinjani, Lombok
2053
2054 1015 WWF. 2002. Resource Economic Valuation of Rinjani Mountain Area. WWF Indonesia Nusa
2055 1016 Tenggara Program, Mataram, Indonesia
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072




Figure 1: Conceptual framework to assess ecosystem services trade-offs (modified from Mouchet et al. 2014)




Figure 2: Map of West Nusa Tenggara province and the remaining forests in Lombok island
(Source: National Institute of Forest Science, Republic of Korea)




Figure 3. Changes in forested area for three 5-year periods (Data source: National Institute of Forest Science, Republic of Korea).






Figure 4: Survey locations (A, B, C, D) and designated forest functions.

Average Surface Runoff (in cubic meters/hour/hectare)

CP Scenario: Change from BAU




FR Scenario: Change from BAU



Average Water Balance (in mm/year)

CP Scenario: Change from BAU



FR Scenario: Change from BAU




Figure 5: Changes in Average Surface Runoff ($\text{m}^3/\text{hour}/\text{ha}$) and Water Balance (mm/year) from recovery of secondary forests in Community Partnership (CP) scenario and recovery of secondary and primary forests in Forest Restoration (FR) scenario




Figure 6: Precipitation records from 1984 to 2014 during dry season: average precipitation from four weather stations near the survey sites around Rinjani Mt and average of six weather stations around the City of Mataram, Lombok, Indonesia (Source: Information Board of Water Resources Province of NTB , 2016)

Table 1: Historical Land Use Changes in Lombok (Unit: ha; Source: Bae at al. 2014)

Land Use Class	1995	2000	2005	2010	Changes 1995-2000	Changes 2000-2005	Changes 2005-2010
Primary forest	54,881	53,140	51,114	51,111	-1,741	-2,025	-4
Secondary forest	105,064	77,452	69,752	67,258	-27,612	-7,700	-2,494
Shrubland	12,767	33,627	42,052	34,419	20,859	8,425	-7,633
All other uses	285,495	293,989	295,289	305,419	8,494	1,300	10,131

Table 2: Carbon stock by land use type (metric ton of carbon/ha \pm standard deviation) (Source: Bae at al. 2014)

		Living vegetation				Dead trees	Litters	Soils
Total	Aboveground							
	Sub-total	Tree	Under growth	Below Ground				
Primary forest	206.6 (±76.66)	109.9	108.6 (±59.89)	1.3 (±1.15)	29.7 (±16.12)	18.3 (±26.05)	1.7 (±1.25)	47.0 (±17.52)
Secondary forest	181.1 (±120.88)	97.8	96.2 (±85.74)	1.6 (±0.99)	26.4 (±23.03)	21.4 (±31.73)	1.8 (±0.84)	33.7 (±13.08)
Shrub land	75.3 (±6.74)	26.5	24.8 (±2.30)	1.7 (±0.98)	7.2 (±0.89)	16.7 (±6.76)	1.6 (±0.43)	23.4 (±3.72)

Table 3: Forest Classification and Permitted Activities (Source: Rosenbarger et al. 2013¹)

Forest classification by function/ Permitted activities ²	Timber Extraction	Cultivating medicinal/decorative plants, fungi, apiculture, swiftlet nests, capturing wildlife, cattle feed	Utilization of environmental services (water flow, ecotourism, biodiversity, environmental protection, carbon absorption and storage)	Extraction of non-timber forest products (rattan, bamboo, honey, resin, fruits, fungi)	Research, science, education, cultivation activities, cultural activities, and limited tourism
Production Forest (A)	Y ³	Y	Y	Y	Y
Protection Forest (B, D ⁴)		Y	Y	Y	Y
Conservation Forest (C)		Y ⁵		Y ⁵	Y

¹ Compiled from: Government Regulation No. 6 of 2007, Minister of Forestry Regulation No. 13 of 2009, Minister of Forestry Regulation No. 37 of 2007, Minister of Forestry Regulation No. 49 of 2008.

² These activities can be legally allowed with permits granted by regent/mayor/governor or minister (depending on area jurisdictions). Although these activities reflect *de facto* uses, two communities in the study area (A and B) do not hold permits.

³ There is no timber concession in the study area.

⁴ The “Community Forest” status of community D means that the forest utilization permit (IUPHKm) was granted to this community for a period of 35 years.

⁵ These activities are not allowed in Conservation Forest, but the community C is in “Traditional Zone”, specially designated for very limited community uses for their livelihoods, including collecting cattle feeds.

Table 4: Potential Land Use Changes under the Business-As-Usual Scenario (ha)

Land Use Class	Present	In 10 years	In 30 years
Primary forest	51,111	51,111	51,111
Secondary forest	67,258	65,462	60,537
Shrubland	34,419	29,030	14,255
All other land uses	305,419	312,604	332,304

Table 5: Potential Land Use Changes under the Community Partnership Scenario (ha)

Land Use Class	Present	In 10 years	In 30 years
Primary forest	51,111	51,111	51,111
Secondary forest	67,258	89,522	105,064
Shrubland	34,419	33,675	12,767
All other land uses	305,419	283,899	289,265

Table 6: Potential Land Use Changes under the Forest Restoration Scenario (ha)

Land Use Class	Present	In 10 years	In 30 years
Primary forest	206.6	52,996	54,881
Secondary forest	67,258	89,522	105,064
Shrubland	34,419	33,675	12,767
All other land uses	305,419	282,014	285,495

Table 7: Land use and Carbon stock change under CP and FR scenarios

Land Use Class	Carbon Stock (metric ton /ha)	Community Partnership scenario (change from BAU) (ha)		Forest Restoration scenario (change from BAU) (ha)	
		in 10 years	in 30 years	in 10 years	in 30 years
Primary forest	206.6	0	0	1,885	3770
Secondary forest	181.1	24,060	44,527	24,060	44,527
Shrubland	75.3	4,645	-1,488	4,645	-1,488
All other land uses	23.4	-28,705	-43,039	-30,590	-46,809
Total carbon stock change (metric tCO ₂ e)		4,035,338	6,944,681	4,380,670	7,635,345

Table 8: Potential Undiscounted Total Market Values of Forest-sequestered Carbon in Lombok (USD millions).

Carbon Price (USD/ metric tCO ₂ e)	Carbon Value (in USD millions)			
	Community Partnership		Forest Restoration	
	10-year	30-year	10-year	30-year
\$5	20.18	34.72	21.90	38.18
\$7.50	30.27	52.09	32.86	57.27
\$10	40.35	69.45	43.81	76.35

Table 9: Level of use (% of respondents reporting collection from forests) and value of forest ESs (USD/household/yr)

Type of service	Forest ESs ¹	Production forest (A)		Protection forest (B)		Conservation forest (C)		Community forest (D)		All forests		% Consumed	
		%	Value	%	Value	%	Value	%	Value	%	Value	%	Sold
Natural NTFP	Bamboo	2	18.52	18	13.35	6	4.23	26	11.25	13	10.83	51	49
	Forage	5	31.11	15	39.21	58	44.39	10	26.67	22	40.49	91	9
	Fern	49	4.22	4	1.63	69	1.48	13	5.04	34	2.86	83	17
	Sub-total	50	8.41	32	20.21	81	27.14	33	14.37	49	18.18		
Agroforest Products	Jackfruit	13	2.79	86	4.23	49	2.47	34	3.31	46	3.47	88	12
	Durian	33	74.80	7	38.27	8	16.89	13	81.63	16	66.46	60	40
	Avocado	17	8.63	29	18.45	43	5.42	3	18.04	23	10.20	44	56
	Mangosteen	3	18.89	0	0.00	0	0.00	1	18.52	1	18.80	44	56
	Melinjo	3	1.44	13	2.55	0	0.00	0	0.00	4	2.31	94	6
	Cacao	28	15.99	14	9.94	0	0.00	0	0.00	11	13.74	8	92
	Coffee	67	262.39	24	50.40	0	0.00	35	63.82	32	171.94	50	50
	Banana	56	14.95	82	15.01	0	0.00	23	13.66	42	14.89	36	64
	Taro	2	14.07	2	2.93	0	0.00	3	4.19	2	7.27	83	17
	Palm	0	0.00	0	0.00	0	0.00	6	918.52	1	918.52	17	83
	Candlenut	0	0.00	16	15.75	5	16.44	3	7.03	6	14.87	31	69
	Other	0	0.00	18	117.18	1	6.73	1	13.46	5	117.45	76	24
	Sub-total	84	142.86	96	49.04	57	14.15	40	103.89	69	77.70		
Timber products	Fuelwood	35	7.17	80	3.59	36	2.99	35	5.92	48	4.56	87	13
	Tools	4	0.74	0	0.00	0	0.00	1	1.85	1	0.96	100	0
	Sub-total	37	6.66	79	3.59	37	2.99	34	5.40	47	4.41		
	All forest ESs²	86	141.49	98	84.98	81	46.25	53	115.63	80	93.46		

¹ No uses were reported for some NTFPs (e.g. langsat, and rattan) and timber products (materials for building and fencing).

² Total % of respondents whose household obtained some values from forest ESs; Mean aggregate value of services obtained from the forest (USD/household/yr).

Table 10: Importance of local forest ESs in future forest management plans by study location

Forest service	Production forest	Protection forest	Conservation forest	Community forest	All respondents
	% of respondents stating that forest service was important				
Natural non-timber forest products	44	26	67	24	40
Agroforest products	92	70	1	86	81
Timber forest products	78	10	1	17	27
Water regulation	96	90	88	90	91

Table 11: Aggregate value of locally provided forest ESs

	Value per year (USD/Household) ¹	Number of affected Households ²	Value per year (million USD)	Undiscounted value over 30 years ³ (million USD)
Production forest	\$121	44,104	\$6.2	\$187
Protection forest	\$83-\$61	84,311	\$7.2-\$9.7	\$241-\$292
Conservation forest	\$38	61,044	\$2.8	\$85
Total		189,460	\$16.2-\$18.8	\$486-\$564

¹ \$121 for Production Forest (\$141 for 86% of the community utilizing forest products); \$83 for Protection Forests (\$85 for 98% of the community utilizing forest products) and \$61 for Community Forests in Protection Forest (\$115 for 53% of the community utilizing forest products) and \$38 for Conservation Forest (\$46 for 81% of the community utilizing forest products)

² Aggregated population of sub-districts near each designated forest function X 51.5% with agriculture as the main occupation based on the 2010 population census.

³ Not accounting for population growth/discounting rate/forest product value change.

Table 12: Changes in value of locally provided forest ESs

	Undiscounted value over 30 years ³ (million USD)	CP scenario ¹		FR scenario ²	
		Forest area changes (%)	Changes in values (million USD)	Forest area changes (%)	Changes in values (million USD)
Production forest	\$187	7.52	\$14.1	8.20	\$15.3
Protection forest	\$241-\$292	18.05	\$43.5-52.5	19.68	\$47.4-57.5
Conservation forest	\$85	12.03	\$10.2	13.12	\$11.2
Total	\$486-\$564	37.6	\$67.8-76.8	41	\$73.9-84.0

¹ 44,527 ha or 37.6% increase in total forest area

² 48,297 ha or 41% increase in total forest area

Table 13: Opportunity costs of carbon sequestration (Value/metric tCO₂e for 30-year)

Profitability of Dryland Agriculture (USD/ha)	Community Partnership (Dryland Agriculture --> Agroforest: 44,527ha)		Forest Restoration (Dryland Agriculture --> Agroforest: 44,527ha & 3,770 ha to primary forest)	
	Undiscounted	NPV with 10% discounting rate	Undiscounted	NPV with 10% discounting rate
\$150	\$0.01	\$0.002	\$0.02	\$0.005
\$250	\$0.13	\$0.04	\$0.13	\$0.04
\$500	\$0.44	\$0.14	\$0.43	\$0.14
\$1,000	\$1.05	\$0.33	\$1.03	\$0.32
\$2,000	\$2.27	\$0.71	\$2.21	\$0.70

*Profitability of Dryland Agriculture/ha – ES value of Forest /ha ((Primary Forest: \$54.58 = \$2.8 million/51,111ha; Secondary/Agroforest: \$144.22 = \$9.7 million/67,258ha); Primary forest = 206.6 metric tCO₂e/ha; Secondary forest = 206.6 metric tCO₂e/ha; Dryland Agriculture = Primary forest = 23.4 metric tCO₂e/ha.